

Heavy metals need assistance: the contribution of nicotianamine to metal circulation throughout the plant and the *Arabidopsis* NAS gene family

Mara Schuler and Petra Bauer *

Department of Biosciences–Plant Biology, Saarland University, Saarbrücken, Germany

Edited by:

Andreas P. M. Weber, University of
Duesseldorf, Germany

Reviewed by:

Elizabeth Pilon-Smits, Colorado State
University, USA

Henk Schat, Vrije Universiteit
Amsterdam, Netherlands

*Correspondence:

Petra Bauer, Department of
Biosciences–Plant Biology, Saarland
University, Campus A2.4, D-66123
Saarbrücken, Germany.
e-mail: p.bauer@mx.uni-saarland.de

Understanding the regulated inter- and intra-cellular metal circulation is one of the challenges in the field of metal homeostasis. Inside organisms metal ions are bound to organic ligands to prevent their uncontrolled reactivity and to increase their solubility. Nicotianamine (NA) is one of the important ligands. This non-proteinogenic amino acid is synthesized by nicotianamine synthase (NAS). NA is involved in mobilization, uptake, transport, storage, and detoxification of metals. Much of the progress in understanding NA function has been achieved by studying mutants with altered nicotianamine levels. Mild and strong *Arabidopsis* mutants impaired in nicotianamine synthesis have been identified and characterized, namely *nas4x-1* and *nas4x-2*. *Arabidopsis thaliana* has four NAS genes. In this review, we summarize the structure and evolution of the NAS genes in the *Arabidopsis* genome. We summarize previous results and present novel evidence that the four NAS genes have partially overlapping functions when plants are exposed to Fe deficiency and nickel supply. We compare the phenotypes of *nas4x-1* and *nas4x-2* and summarize the functions of NAS genes and NA as deduced from the studies of mutant phenotypes.

Keywords: nicotianamine, metal binding, chelation, gene family, multiple mutant

INTRODUCTION

Iron (Fe) and copper (Cu) are essential elements for all living organisms because of their unique property of being able to catalyze oxidation/reduction reactions. Conversely, an excess of Fe, especially Fe^{2+} , is detrimental since it catalyzes the production of reactive oxygen species (ROS) in the Fenton reaction (Fenton, 1894; Hell and Stephan, 2003). For this reason, free metal ions are not likely to exist in large amounts in cells. Indeed, Fe and other metals are mainly present in stable complexes with organic ligands or inorganic phosphates (Haydon and Cobbett, 2007).

Nicotianamine (NA) is one of the most investigated metal chelator molecules in plants. NA is a non-proteinogenic amino acid and it results from the enzymatic condensation of three S-adenosyl methionine molecules (SAM) catalyzed by nicotianamine synthase (NAS; Herbig et al., 1999; Ling et al., 1999; Takahashi et al., 1999). NA is able to form stable complexes with Mn^{2+} , Fe^{2+} , Co^{2+} , Zn^{2+} , Ni^{2+} , and Cu^{2+} (Benes et al., 1983; Anderegg and Ripberger, 1989). Moreover, NA has a high capacity to chelate Fe^{3+} (von Wiren et al., 1999; Weber et al., 2006; Rellán-Alvarez et al., 2008). For all the metals considered, the stability of the NA–metal complexes had its maximum at pH values above 6.5, suggesting that NA is more likely a symplastic chelator of metals and therefore would bind metals predominately within cells and the phloem (von Wiren et al., 1999). Cu^{2+} is an exception among the essential metals, since the Cu^{2+} –NA complex is very stable in mild acidic conditions, which is a strong argument in favor of

the possible occurrence of Cu^{2+} –NA complexes in the apoplastic environment as prevailing in the xylem sap (von Wiren et al., 1999). Nicotianamine can be transported to the various organs and tissues via oligopeptide transporters, such as yellowstripe1-like (YSL) proteins (Curie et al., 2001, 2009). Rice ENA1 and ENA2 transporters were just recently described to mediate NA export from cells (Nozoye et al., 2011).

Studies of solanaceous and graminaceous plants as well as of hyperaccumulators showed that NA functions in long-distance transport of Cu (Pich and Scholz, 1996), short-distance and intracellular transport of Fe (Becker et al., 1995; Curie and Briat, 2003), plant reproduction (Takahashi et al., 2006), detoxification of heavy metals like Ni (Douchkov et al., 2005; Kim et al., 2005; Pianelli et al., 2005; Mari et al., 2006; Ouerdane et al., 2006; van de Mortel et al., 2006; Callahan et al., 2007) and Zn (Becher et al., 2004; Weber et al., 2004; Talke et al., 2006; van de Mortel et al., 2006), and in grasses as a precursor in the biosynthesis of phytosiderophores (Mori and Nishizawa, 1987). Several studies suggested that NA could be involved in iron mobilization and accumulation in plant roots and seeds (Douchkov et al., 2001, 2005; Cheng et al., 2007; Lee et al., 2009). NA is beneficial for increased bioavailability of Fe in foods (Maurer et al., 2010; Zheng et al., 2010).

Taken together, NA is a key compound of metal homeostasis in plants contributing to mobilization, uptake, transport, storage, and detoxification of metals. Since NA was found to be an important biofortification factor for essential nutrients like Fe and Zn in edible portions of crop plants (Zheng et al., 2010), further knowledge about the functions of NA and the

Abbreviations: NA, nicotianamine; NAS, nicotianamine synthase.


characterization of the essential genes for its production are of high relevance.

NAS GENE FAMILY OF A. THALIANA

The *Arabidopsis* system provides all tools that allow combining genetic studies with physiological analyzes and global gene expression experiments. This species was therefore utilized to investigate NA function. While not all plant species have multiple *NAS* genes, the *Arabidopsis* genome harbors a *NAS* gene family comprising four members (Bauer et al., 2004). *NAS1* and *NAS2* are located on chromosome V, while *NAS3* and *NAS4* are located on chromosome I (Figure 1A). Multiple alignment (CLUSTALW) showed a close relation between *NAS* genes located on the same chromosome

with more than 80% identity while alignment of genes belonging to separate chromosomes showed an identity of about 70% (Bauer et al., 2004). Gene mapping between *Solanum esculentum* and *A. thaliana* suggested that the four *Arabidopsis* genes, as well as the single tomato *NAS* gene originated from a common ancestor *NAS* gene. This finding is in agreement with a first genome duplication event in the evolution of *Arabidopsis*, followed later by two independent duplication events (Bauer et al., 2004).

T-DNA insertion lines of all four *NAS* genes were identified and crossed to each other (Figure 1B; primer sequences for genotyping are found in Figure 1C). Under regular growth conditions single, double, and triple mutants did not show any obvious phenotypes suggesting functional redundancy (Klatte et al., 2009).



Single mutants had similar NA contents as wild type. Triple *nas* mutants had NA levels that were reduced to 30–40% of wild type levels (Klatte et al., 2009). Since NA can be transported short and long-distance in plants, severe metal homeostasis phenotypes are not expected in the presence of a functional *NAS* gene. Interestingly, however, upon exposure to modestly toxic Ni supply, *nas4-1* had a more chlorotic phenotype than *nas3-1*, while *nas1-1* and *nas2-1* had mild phenotypes like the wild type (Klatte et al., 2009). With increasing number of *NAS* knockout alleles, the NA contents decreased in the mutants while the severity of the leaf chlorosis was enhanced (Klatte et al., 2009). This suggests that NA contents correlate with Ni tolerance. Here, we show the seedling growth responses of single and multiple mutants in response to Fe deficiency. We found that all single *nas* mutants tested had a stronger leaf chlorosis than wild type plants upon Fe deficiency (Figure 2A). *nas4-1* Mutants had the strongest leaf chlorosis among the tested single mutants (Figure 2A). It can therefore be concluded that the *NAS* gene functions are partially non-overlapping. Perhaps the location of NA production is important. Alternatively, the *NAS* isoforms might have different enzyme activities, perhaps under specific conditions like Fe deficiency and Ni supply.

Partial non-redundancy is further confirmed by the fact that the *NAS* genes are differentially regulated in plants (Bauer et al., 2004; Klatte et al., 2009; summarized in Figure 2B; primers for gene expression analysis in Figure 1C). *NAS1*, *NAS2*, and *NAS4* were found expressed in roots, where they were induced by Ni supply. *NAS2* was also up-regulated by Fe deficiency. *NAS1* and *NAS4* were expressed in leaves, and *NAS4* could be induced by Fe deficiency

and Ni in leaves. *NAS2* was not expressed in leaves. On the other hand, *NAS3* was expressed in leaves where it was repressed by Fe deficiency but strongly induced by Ni supply. In flowers, *NAS3* was expressed in sepals and petals while *NAS1*, *NAS2*, and *NAS4* were not expressed (Schuler et al., in preparation).

Taken together, *NAS* genes evolved as a gene family in *Arabidopsis* where they acquired overlapping and specific functions in metal homeostasis as well as differential gene regulation in response to metals. *NAS3* seems important for leaf and flower nicotianamine production upon Fe supply as well as Ni tolerance, while *NAS4* was more important for Fe deficiency in leaves and perhaps in roots. *NAS2* might be especially relevant for Fe deficiency responses in roots.

PHYSIOLOGICAL ANALYSIS OF QUADRUPLE *NAS* MUTANTS

Mutant analysis showed that all four *NAS* genes are functional, so that quadruple mutant analysis was needed to study NA function. The leaf chlorosis phenotypes of quadruple *nas* mutants were more severe than those of single mutants. Quadruple *nas1-1 nas2-1 nas3-1 nas4-1* mutants (termed *nas4x-1*) were found to have a stronger reduction of NA levels than all triple mutant combinations analyzed, namely to approximately 15% in vegetative leaves and 30% in seeds compared to wild type (Klatte et al., 2009). While *nas4x-1* plants had a residual NA level in leaves at the vegetative stage, this was not the case in the reproductive stage in leaves. Full loss of function *nas1-1 nas2-2 nas3-1 nas4-1* mutants (termed *nas4x-2*) did not contain any NA (Klatte et al., 2009). The morphological phenotypes of *nas4x-1* and *nas4x-2* were compared (summarized in Table 1).




Table 1 | Comparison of *nas4x-1* and *nas4x-2*.

	<i>nas4x-1</i>	<i>nas4x-2*</i>	Reference
Genotype	<i>nas1-1 nas2-1 nas3-1 nas4-1</i>	<i>nas1-1 nas2-2 nas3-1 nas4-1</i>	Klatte et al. (2009)
Full-length <i>NAS</i> transcripts	<i>NAS2</i>	none	Klatte et al. (2009)
NA content, versus WT	10% in leaves, vegetative stage; 0% in leaves, reproductive stage; 40% in seeds	0% in all tested leaves; no seeds produced	Klatte et al. (2009)
Interveinal leaf chlorosis	Starts during reproductive phase	Strong during vegetative and reproductive phase; more severe in young leaves than in older leaves	Klatte et al. (2009)
Numerous small rosette leaves	Yes	Yes	Klatte et al. (2009)
Flowers	Normal	Sterile	Klatte et al. (2009)
Flowering time and senescence	Normal	Delayed	Klatte et al. (2009)
Global gene expression	Categories affected: metal homeostasis, biotic stress responses, leaf photosystem and root carbohydrate metabolism	n.a.	Schuler et al. (2011)

*A more detailed investigation of *nas4x-2* phenotypes will be presented in Schuler et al. (in preparation); n.a. = not analyzed.

nas4x-1 plants appeared nearly normal during the vegetative stage, unless they were grown under Fe deficiency or Ni supply. However, *nas4x-1* plants showed an interveinal leaf chlorosis upon transition to the reproductive growth stage, and Fe contents were increased in leaves at this stage compared to wild type (Klatte et al., 2009). Mobilization of Fe by *nas4x-1* roots was up-regulated at this stage which accounts for the increased Fe contents (Klatte et al., 2009). Presumably, intercostal leaf areas with mesophyll cells did not acquire Fe in sufficient amounts and may have emitted a long-distance Fe deficiency signal that stimulated root Fe uptake. *nas4x-1* plants are still fertile, yet flowers and seeds were found to contain less Fe than in the wild type (Klatte et al., 2009). *nas4x-1* mutant plants are valuable models to study NA function in late phases of plant development. These plants also served to perform a transcriptome analysis of roots and leaves upon Fe supply and Fe deficiency (Schuler et al., 2011). A comparison with the wild type transcriptomes confirmed that *nas4x-1* was affected in metal homeostasis since a high number of genes of this category was hit by differential expression. Besides this category, the mutant was also affected in biotic stress responses, leaf photosystem organization, and root carbohydrate metabolism (Schuler et al., 2011). Significantly more genes of these four biological categories were affected by differential expression between mutant and wild type compared to all genes analyzed in the microarray study. A change of expression of genes from these categories can be explained as an adaptation response to altered Fe levels.

Nas4x-2, on the other hand, is a severely affected mutant. Leaf chlorosis started during the vegetative phase (Klatte et al., 2009).

Closer inspection of this mutant showed that NA was involved in the long-distance transport of Fe to young leaves presumably using the phloem, while older leaves received Fe from citrate-mediated transport in the xylem (Schuler et al., in preparation). The leaf chlorosis was due to Fe accumulation in the vascular system suggesting that NA is involved in lateral transport of Fe from vascular tissues to mesophyll (Schuler et al., in preparation). Furthermore, *nas4x-2* mutants were affected in pollination (Schuler et al., in preparation).

CONCLUDING REMARKS

Arabidopsis served as a model for the study of nicotianamine function in plants. The split of a *NAS* locus to four *NAS* genes in *Arabidopsis* resulted in a partial non-redundant specialization of *NAS* gene functions. These are conferred at least partly by differential gene expression of the *NAS* genes in response to developmental cues, tissue specificity, and metals. It has not been investigated yet whether the enzyme activities of the *NAS* isoforms are differentially regulated by metal supply. Using the mild and severe *nas4x-1* and *nas4x-2* mutants novel nicotianamine functions were uncovered, such as seed Fe loading, long-distance Fe transport to leaves, short-distance transport from vascular tissues to mesophyll, and in pollination.

ACKNOWLEDGMENTS

This work was funded by the DFG grants (Ba 1610/4–1, 4–4, 6–1) within the framework of the *Arabidopsis* Functional Genomics Network (AFGN).

REFERENCES

Anderegg, G., and Ripberger, H. (1989). Correlation between metal complex formation and biological activity of nicotianamine analogues. *J. Chem. Soc. Chem. Commun.* 647–650.

Bauer, P., Thiel, T., Klatte, M., Bereczky, Z., Brumbarova, T., Hell, R., and Grosse, I. (2004). Analysis of sequence, map position, and gene expression reveals conserved essential genes for iron uptake in *Arabidopsis* and tomato. *Plant Physiol.* 136, 4169–4183.

Becher, M., Talke, I. N., Krall, L., and Krämer, U. (2004). Cross-species microarray transcript profiling reveals high constitutive expression of metal homeostasis genes in shoots of the zinc hyperaccumulator *Arabidopsis halleri*. *Plant J.* 37, 251–268.

Becker, R., Fritz, E., and Manteuffel, R. (1995). Subcellular localization and characterization of excessive iron in the nicotianamine-less tomato mutant chloronerva. *Plant Physiol.* 108, 269–275.

Benes, I., Schreiber, K., Rippberger, H., and Kircheiss, A. (1983). Metal complex formation of nicotianamine, a possible phytosiderophore. *Experientia* 39, 261–262.

Callahan, D. L., Kolev, S. D., O’Hair, R. A. J., Salt, D. E., and Baker, A. J. M. (2007). Relationships of nicotianamine and other amino acids with nickel, zinc and iron in *Thlaspi* hyperaccumulators. *New Phytol.* 176, 836–848.

Cheng, L. J., Wang, F., Shou, H. X., Huang, F. L., Zheng, L. Q., He, F., Li, J. H., Zhao, F. J., Ueno, D., Ma, J. F., and Wu, P. (2007). Mutation

- in nicotianamine aminotransferase stimulated the Fe(II) acquisition system and led to iron accumulation in rice. *Plant Physiol.* 145, 1647–1657.
- Curie, C., and Briat, J.-F. (2003). Iron transport and signaling in plants. *Annu. Rev. Plant Biol.* 54, 183–206.
- Curie, C., Cassin, G., Couch, D., Divol, F., Higuchi, K., Le Jean, M., Misson, J., Schikora, A., Czernic, P., and Mari, S. (2009). Metal movement within the plant: contribution of nicotianamine and yellow stripe 1-like transporters. *Ann. Bot.* 103, 1–11.
- Curie, C., Panaviene, Z., Loulergue, C., Dellaporta, S. L., Briat, J. F., and Walker, E. L. (2001). Maize yellow stripe1 encodes a membrane protein directly involved in Fe(III) uptake. *Nature* 409, 346–349.
- Douchkov, D., Gryczka, C., Stephan, U. W., Hell, R., and Baumlein, H. (2005). Ectopic expression of nicotianamine synthase genes results in improved iron accumulation and increased nickel tolerance in transgenic tobacco. *Plant Cell Environ.* 28, 365–374.
- Douchkov, D., Hell, R., Stephan, U. W., and Baumlein, H. (2001). Increased iron efficiency in transgenic plants due to ectopic expression of nicotianamine synthase. *Plant Nutr.* 92, 54–55.
- Fenton, H. J. H. (1894). Oxidation of tartaric acid in presence of iron. *J. Chem. Soc. Trans.* 65, 899–910.
- Haydon, M. J., and Cobbett, C. S. (2007). Transporters of ligands for essential metal ions in plants. *New Phytol.* 174, 499–506.
- Hell, R., and Stephan, U. W. (2003). Iron uptake, trafficking and homeostasis in plants. *Planta* 216, 541–551.
- Herbik, A., Koch, G., Mock, H. P., Dushkov, D., Czihal, A., Thielmann, J., Stephan, U. W., and Baumlein, H. (1999). Isolation, characterization and cDNA cloning of nicotianamine synthase from barley – a key enzyme for iron homeostasis in plants. *Eur. J. Biochem.* 265, 231–239.
- Jakoby, M., Wang, H. Y., Reidt, W., Weishaar, B., and Bauer, P. (2004). FRU (BHLH029) is required for induction of iron mobilization genes in *Arabidopsis thaliana*. *FEBS Lett.* 577, 528–534.
- Kim, S., Takahashi, M., Higuchi, K., Tsunoda, K., Nakanishi, H., Yoshimura, E., Mori, S., and Nishizawa, N. K. (2005). Increased nicotianamine biosynthesis confers enhanced tolerance of high levels of metals, in particular nickel, to plants. *Plant Cell Physiol.* 46, 1809–1818.
- Klatte, M., Schuler, M., Wirtz, M., Fink-Straube, C., Hell, R., and Bauer, P. (2009). The analysis of *Arabidopsis* nicotianamine synthase mutants reveals functions for nicotianamine in seed iron loading and iron deficiency responses. *Plant Physiol.* 150, 257–271.
- Klatte, M., and Bauer, P. (2009). Accurate real-time reverse transcription quantitative PCR. *Methods Mol. Biol.* 479, 61–77.
- Lee, S., Jeon, U. S., Lee, S. J., Kim, Y.-K., Persson, D. P., Husted, S., Schjorring, J. K., Kakei, Y., Masuda, H., Nishizawa, N. K., and An, G. (2009). Iron fortification of rice seeds through activation of the nicotianamine synthase gene. *Proc. Natl. Acad. Sci. U.S.A.* 106, 22014–22019.
- Ling, H., Koch, Q., Baumlein, G. H., and Ganai, M. W. (1999). Map-based cloning of chloronerva, a gene involved in iron uptake of higher plants encoding nicotianamine synthase. *Proc. Natl. Acad. Sci. U.S.A.* 96, 7098–7103.
- Mari, S., Gendreau, D., Pianelli, K., Ouerdane, L., Lobinski, R., Briat, J. F., Lebrun, M., and Czernic, P. (2006). Root-to-shoot long distance circulation of nicotianamine and nicotianamine-nickel chelates in the metal hyperaccumulator *Thlaspi caerulescens*. *J. Exp. Bot.* 57, 4111–4122.
- Maurer, F., Daum, N., Schaefer, U. F., Lehr, C. M., and Bauer, P. (2010). Plant genetic factors for iron homeostasis affect iron bioavailability in Caco-2 cells. *Food Res. Int.* 43, 1661–1665.
- Mori, S., and Nishizawa, N. K. (1987). Methionine as a dominant precursor of phytosiderophores in Gramineae plants. *Plant Cell Physiol.* 28, 1081–1092.
- Nozoye, T., Nagasaka, S., Kobayashi, T., Takahashi, M., Sato, Y., Sato, Y., Uozumi, N., Nakanishi, H., and Nishizawa, N. K. (2011). Phytosiderophore efflux transporters are crucial for iron acquisition in graminaceous plants. *J. Biol. Chem.* 286, 5446–5454.
- Ouerdane, L., Mari, S., Czernic, P., Lebrun, M., and Lobinski, R. (2006). Speciation of non-covalent nickel species in plant tissue extracts by electrospray Q-TOFMS/MS after their isolation by 2D size exclusion-hydrophilic interaction LC (SEC-HILIC) monitored by ICP-MS. *J. Anal. At. Spectrom.* 21, 676.
- Pianelli, K., Mari, S., Marquès, L., Lebrun, M., and Czernic, P. (2005). Nicotianamine over-accumulation confers resistance to nickel in *Arabidopsis thaliana*. *Transgenic Res.* 14, 739–748.
- Pich, A., and Scholz, G. (1996). Translocation of copper and other micronutrients in tomato plants (*Lycopersicon esculentum* Mill.): nicotianamine-stimulated copper transport in the xylem. *J. Exp. Bot.* 47, 41–47.
- Relan-Alvarez, R., AbadiÁa, J., and Alvarez-Fernandez, A. (2008). Formation of metal-nicotianamine complexes as affected by pH, ligand exchange with citrate and metal exchange. A study by electrospray ionization time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* 22, 1553–1562.
- Schuler, M., Keller, A., Backes, C., Phillipar, K., Lenhof, H. P., and Bauer, P. (2011). Transcriptome analysis by GeneTrail revealed regulation of functional categories in response to alterations of iron homeostasis in *Arabidopsis thaliana*. *BMC Plant Biol.* 11, 87. doi:10.1186/1471-2229-11-87
- Takahashi, M., Inoue, H., Ishimaru, Y., Nakanishi, H., Mori, S., and Nishizawa, N. (2006). The role of nicotianamine and mugineic acid in metal transport at reproductive stage. *Plant Cell Physiol.* 47, S230–S230.
- Takahashi, M., Yamaguchi, H., Nakanishi, H., Shioiri, T., Nishizawa, N. K., and Mori, S. (1999). Cloning two genes for nicotianamine aminotransferase, a critical enzyme in iron acquisition (strategy II) in graminaceous plants. *Plant Physiol.* 121, 947–956.
- Talke, I. N., Hanikenne, M., and Kramer, U. (2006). Zinc-dependent global transcriptional control, transcriptional deregulation, and higher gene copy number for genes in metal homeostasis of the hyper accumulator *Arabidopsis halleri*. *Plant Physiol.* 142, 148–167.
- van de Mortel, J. E., Villanueva, L. A., Schat, H., Kwekkeboom, J., Coughlan, S., Moerland, P. D., van Themaat, E. V. L., Koornneef, M., and Aarts, M. G. M. (2006). Large expression differences in genes for iron and zinc homeostasis, stress response, and lignin biosynthesis distinguish roots of *Arabidopsis thaliana* and the related metal hyperaccumulator *Thlaspi caerulescens*. *Plant Physiol.* 142, 1127–1147.
- von Wiren, N., Klair, S., Bansal, S., Briat, J. F., Khodr, H., Shioiri, T., Leigh, R. A., and Hider, R. C. (1999). Nicotianamine chelates both Fe-III and Fe-II. Implications for metal transport in plants. *Plant Physiol.* 119, 1107–1114.
- Weber, G., von Wiren, N., and Hayen, H. (2006). Analysis of iron (II)/iron(III) phytosiderophore complexes by nano-electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry. *Rapid Commun. Mass Spectrom.* 20, 973–980.
- Weber, M., Harada, E., Vess, C., Roepenack-Lahaye, E. V., and Clemens, S. (2004). Comparative microarray analysis of *Arabidopsis thaliana* and *Arabidopsis halleri* roots identifies nicotianamine synthase, a ZIP transporter and other genes as potential metal hyper accumulating factors. *Plant J.* 37, 269–281.
- Zheng, L., Cheng, Z., Ai, C., Jiang, X., Bei, X., Zheng, Y., Glahn, R. P., Welch, R. M., Miller, D. D., Lei, X. G., and Shou, H. (2010). Nicotianamine, a novel enhancer of rice iron bioavailability to humans. *PLoS ONE* 5, e10190. doi:10.1371/journal.pone.0010190

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 31 August 2011; accepted: 10 October 2011; published online: 22 November 2011.

Citation: Schuler M and Bauer P (2011) Heavy metals need assistance: the contribution of nicotianamine to metal circulation throughout the plant and the *Arabidopsis* NAS gene family. *Front. Plant Sci.* 2:69. doi: 10.3389/fpls.2011.00069
This article was submitted to *Frontiers in Plant Physiology*, a specialty of *Frontiers in Plant Science*.

Copyright © 2011 Schuler and Bauer. This is an open-access article subject to a non-exclusive license between the authors and Frontiers Media SA, which permits use, distribution and reproduction in other forums, provided the original authors and source are credited and other Frontiers conditions are complied with.